

Aquatic Invertebrate Monitoring at Effigy Mounds National Monument, 2008

Natural Resource Data Series NPS/HTLN//NRDS—2010/071



Aquatic Invertebrate Monitoring at Effigy Mounds National Monument, 2008

Natural Resource Data Series NPS/HTLN//NRDS—2010/071

David E. Bowles and Hope R. Dodd National Park Service The Heartland I&M Network Wilson's Creek National Battlefield 6424 West Farm Road 182 Republic, MO 65738

Jessica A. Luraas Department of Biology Missouri State University 901 South National Avenue Springfield, MO 65897

August 2010

U.S. Department of the Interior National Park Service Natural Resource Program Center Fort Collins, Colorado The National Park Service, Natural Resource Program Center publishes a range of reports that address natural resource topics of interest and applicability to a broad audience in the National Park Service and others in natural resource management, including scientists, conservation and environmental constituencies, and the public.

The Natural Resource Data Series is intended for timely release of basic data sets and data summaries. Care has been taken to assure accuracy of raw data values, but a thorough analysis aninterpretation of the data has not been completed. Consequently, the initial analyses of data in this report are provisional and subject to change.

All manuscripts in the series receive the appropriate level of peer review to ensure that the information is scientifically credible, technically accurate, appropriately written for the intended audience, and designed and published in a professional manner. This report received informal peer review by subject-matter experts who were not directly involved in the collection, analysis, or reporting of the data.

Views, statements, findings, conclusions, recommendations, and data in this report do not necessarily reflect views and policies of the National Park Service, U.S. Department of the Interior. Mention of trade names or commercial products does not constitute endorsement or recommendation for use by the U.S. Government.

This report is available from http://science.nature.nps.gov/im/units/htln/ and the Natural Resource Publications Management website (http://www.nature.nps.gov/publications/NRPM/).

Please cite this publication as:

Bowles D. E., H. R. Dodd, and J. A. Luraas. 2010. Aquatic invertebrate monitoring at Effigy Mounds National Monument, 2008. Natural Resource Data Series NPS/HTLN/NRDS—2010/071. National Park Service, Fort Collins, Colorado.

Contents

	Page
Tables	iv
Abstract	v
Acknowledgments	vi
Introduction	1
Methods	1
Results	3
Discussion	6
Literature cited	7
Appendix	8

Tables

Pag	е
Table 1. Summary statistics for invertebrate samples collected from Dousman Creek, Effigy Mounds National Monument, 2008.	4
Table 2 . Water quality data for Dousman Creek, Effigy Mounds National Monument, 2008. Data were collected hourly with calibrated data loggers.	
Table 3. Statistical ranges of water quality parameters sampled at 98 candidate reference sites from 1994-1998 (from Wilton 2004).	
Table 4. Summary statistics for habitat variables associated with benthic samples collected from Dousman Creek, Effigy Mounds National Monument, 2008.	

Abstract

The National Park Service (NPS) began monitoring water quality and invertebrate community structure in Dousman Creek at Effigy Mounds National Monument (EFMO) in 2008. Dousman Creek is a Class "B" coldwater stream that supports a trout population and is listed among the Outstanding Iowa Waters. The purpose of this report is to summarize baseline aquatic invertebrate monitoring data collected during July 2008. Data are currently insufficient to fully characterize the integrity of Dousman Creek. Preliminary indications based on invertebrate community structure, water quality and physical habitat suggests Dousman Creek is largely unimpaired. All benthic samples were dominated by a moderately intolerant mayfly (Pseudocloeon) and Chironomidae were poorly represented in samples. Because large numbers of Chironomidae are often associated with disturbance, their low density in Dousman Creek is particularly notable. Taxa richness and EPT richness were lower than expected for this outstanding stream, but the reason for this is unknown. Several intolerant and uncommon taxa were represented in samples. Water quality data also suggest this stream is not disturbed. Future invertebrate sampling events at Dousman Creek should include a composited multihabitat sample that will allow investigators to calculate the IOWATER Advanced Benthic Macroinvertebrate Indexing Methods. Using this index to evaluate future benthic collections from Dousman Creek will allow site managers to directly compare its invertebrate community structure relative to Iowa reference streams in addition to assessing long-term trend in this resource.

Acknowledgments

We thank Jan Hinsey, Ryan Green, Myranda Clark, Jennifer Haack, Lloyd Morrison, and Rodney Rovang for their assistance on this project. Brandy Bregthold (Missouri Department of Natural Resources), and Sam Testa (USDA National Sedimentation Lab) kindly served as peer reviewers on an earlier draft of this report.

Introduction

Aquatic invertebrates are an important biological tool for understanding and detecting changes in stream ecosystem integrity, and they can be used to reflect cumulative impacts that cannot otherwise be detected through traditional water quality monitoring. The National Park Service (NPS) began monitoring water quality and invertebrate community structure in Dousman Creek at Effigy Mounds National Monument (EFMO) in 2008 (Bowles et al. 2008). Dousman Creek is a Class "B" coldwater stream that supports a trout population and is listed among the Outstanding Iowa Waters (Iowa Department of Natural Resources, 2010a). The monitoring objectives of this study, as described by DeBacker *et al.* (2005), are: 1) determine the status and trends of invertebrate species diversity, abundance, and community metrics, and 2) relate the invertebrate community to overall water quality through quantification of metrics related to taxa richness, abundance, diversity, and region-specific multi-metric indices as indicators of water quality and habitat condition. The purpose of this report is to summarize baseline aquatic invertebrate monitoring data collected during July 2008.

Methods

Methods and procedures used in this report follow Bowles *et al.* (2008), Monitoring Protocol for Aquatic Invertebrates of Small Streams in the Heartland Inventory & Monitoring Network. Samples were collected at one reach of Dousman Creek on July 7, 2008 (Figure 1).

Three successive riffles were sampled with three benthic invertebrate samples collected at each riffle, resulting in nine samples. A Surber stream bottom sampler (500 µm mesh, 0.09 m²) was used to collect the samples. Samples were sorted in the laboratory following a subsampling routine described in Bowles *et al.* (2008). Taxa were identified to the lowest practical taxonomic level (usually genus) and counted. Metrics calculated for each sample included genus richness, Shannon diversity index, EPT (Ephemeroptera, Plecoptera, Trichoptera) richness, EPT ratio (EPT density/(EPT density + Chironomidae density)), genus evenness (where 0 = minimum evenness, 1 = maximum evenness), and Hilsenhoff biotic index (HBI) (Bowles et al. 2008). For details on calculating and interpreting metrics used in this report refer to Bowles et al. (2008). The preferred response of all metrics is an increase except for HBI where smaller values are the desired response. An increase in HBI is undesired because that would reflect increasing tolerance of the community to disturbance.

For each sample, current velocity (meters/second) and depth (cm) were recorded directly in front of the sampling net frame. Qualitative habitat variables (embeddedness, periphyton, filamentous algae, aquatic vegetation, deposition, and organic material) were estimated within the sampling net frame as percentage categories (0, <10, 10-40, 40-75, >75). Habitat data were analyzed as midpoints of each category. Dominant substrate size from the area within the sampling net frame was visually assessed using the Wentworth scale (Wentworth 1922). Stream discharge was measured upstream of the sampled riffles. Seven water quality readings were recorded hourly using a calibrated YSI 6920 data logger. The water quality and habitat data presented in this report represent only a snap-shot of the broad temporal range of conditions and should be cautiously interpreted. They are intended to describe the prevailing conditions that influence the structure of invertebrate communities, and they may help explain variability between samples,

but they should not be used as an analytical tool in the strictest sense (Bowles *et al.* 2008). Due to the limitations of using water quality data obtained with data loggers, the invertebrate community is used here as a surrogate of the long-term water quality condition of Dousman Creek.

The intent of this study is to monitor community condition within a site over time and not make comparisons within the site (Bowles et al. 2008). Because sample size presently is small (n=1 year), a rigorous statistical analysis of the data cannot be accomplished and only summary statistics are presented in this report.




Figure 1. Map showing the approximate lower sampling reach boundary (yellow dot) for Dousman Creek, Effigy Mounds National Monument.

Results

Genus and EPT richness values for Dousman Creek were lower than those expected for regional reference streams (Wilton 2004). Mean genus richness across riffles was 10.11 (range=9.0-11.3) and roughly one fourth of those taxa were EPT (Table 1). Among taxa represented in samples, the mean tolerance value (TV) was 5.4 (range 0-10 with 10 being most tolerant). Roughly onefourth of all taxa collected were sensitive having tolerance values of 3 or less. Sensitive taxa present in samples included the caddisfly (Trichoptera) genera Brachycentrus, Glossomsoma and Neophylax, the riffle beetle Optioservus (Coleoptera), and the cranefly larva Dicranota (Diptera) (Appendix). Although EPT taxa were poorly represented among samples, the dominant taxon among all taxa in every sample was the mayfly (Ephemeroptera) genus *Pseudocloeon* (TV = 4.4). Other common taxa represented in samples were black fly larvae (Diptera: Simulium), and the Crustaceans Lirceus (Isopoda) and Gammarus (Amphipoda), with tolerance values of 4.4, 7.7., and 6.9, respectively. The relatively high EPT ratio for Dousman Creek (0.89) indicates that the dipteran family Chironomidae did not represent a substantial portion of the benthic community among samples. The percentage composition of Chironomidae in the total benthic community was low and ranged from 3.76% to19.11% (mean=8.18%). Shannon index among samples was relatively low ranging from 1.12 to 1.48 (mean=1.26). Shannon's index accounts for both abundance and evenness of the species present and index values are more robust when all taxa in a sample are equally abundant or have high evenness. The dominance of a few taxa among samples, particularly *Pseudocloen*, is reflected in the low Shannon Index values reported here. Genus evenness ranged from 0.49-0.61 (mean=0.55). HBI was relatively low among samples with scores ranging from 4.60-4.85 (mean=4.70). Because HBI scores can range from 0 to 10 with ten indicating the most disturbance, the relatively low HBI scores for Dousman Creek indicate the water quality condition for that stream is good.

All water quality parameters measured in this study compared favorably to Iowa surface water standards and to regional reference streams (Tables 2 and 3). The mean water temperature of about 15 °C indicates Dousman Creek is a coldwater stream (Wilton 2004). The Iowa surface water quality standard for dissolved oxygen is that the minimum dissolved oxygen levels shall not be less than 5.0 mg/l during any 24-hour period (Iowa Department of Natural Resources 2010b). Dissolved oxygen levels in Dousman Creek clearly surpassed this standard. Stream discharge at the time of sampling was 0.42 m³.

Habitat among riffles was uniform (Table 4). Riffles were shallow (mean= 16.89 cm) with moderate current velocity (mean=0.54 m/sec). Substrate among riffles consisted primarily of small to large pebble with size ranging from 14.33-14.67 (mean= 14.44, or 32-45 mm) on the Wentworth scale (Bowles 2008). Embeddedness and deposition were relatively low (means= 25% and 26.39%, respectively). Mean periphyton was 18.33% and did not exceed 25%. No aquatic vegetation or filamentous algae was present in samples. Percent organic material in samples also was low ranging from 5-25% (mean 13.89%).

Table 1. Summary statistics for invertebrate samples collected from Dousman Creek, Effigy Mounds National Monument, 2008.

Statistic	НВІ	Genus Richness	Shannon's Index	Genus Evenness	EPT Richness	EPT Ratio
Mean	4.70	10.00	1.26	0.55	2.11	0.89
Standard Error	0.07	0.69	0.11	0.03	0.11	0.02
Minimum	4.60	9.00	1.11	0.49	2.00	0.86
Maximum	4.85	11.33	1.48	0.61	2.33	0.93
N	9	9	9	9	9	9

Table 2. Water quality data for Dousman Creek, Effigy Mounds National Monument, 2008. Data were collected hourly with calibrated data loggers.

Water Quality Statistic	Temperature (°C)	Specific Conductance (µS/cm)	Dissolved Oxygen (mg/liter)	рН	Turbidity (NTU)
Mean	15.21	562.43	10.52	8.05	3.2
Median	15.40	562.00	10.39	8.07	2.9
Standard Deviation	1.38	2.99	0.44	0.05	1.16
Minimum	12.97	560.00	10.03	7.94	2.00
Maximum	17.00	569.00	11.33	8.08	5.10
Count	7	7	7	7	6

Table 3. Statistical ranges of water quality parameters sampled at 98 candidate reference sites from 1994-1998 (from Wilton 2004).

Parameter	Minimum	25th Percentile	50th Percentile	75th Percentile	Maximum
Temperature (°C)	8	14.6	18.6	21	26.8
Dissolved Oxygen (mg/l)	4.7	7.7	8.4	9.5	12.6
рН	6.5	7.4	7.7	8	8.6
Conductivity (µmhos/cm)	340	518	625	733	1200
Turbidity (NTU)	1	8	16	26	80

Table 4. Summary statistics for habitat variables associated with benthic samples collected from Dousman Creek, Effigy Mounds National Monument, 2008.

Habitat Parameter	Mean	Standard Error	Minimum	Maximum	N
Depth (cm)	16.89	3.39	10.67	22.33	9
Current velocity (m/sec)	0.54	0.11	0.33	0.69	9
Substrate (Wentworth Scale)	14.44	0.11	14.33	14.67	9
Embeddedness (%)	25	0	25	25	9
Vegetation (%)	0	0	0	0	9
Filamentous algae (%)	0	0	0	0	9
Periphyton (%)	18.33	6.67	5	25	9
Deposition (%)	26.39	5.09	18.33	35.83	9
Organics (%)	13.89	5.88	5	25	9

Discussion

Data are currently insufficient to fully characterize the integrity of Dousman Creek. Preliminary indications based on invertebrate community structure, water quality and physical habitat suggests Dousman Creek is largely unimpaired. All benthic samples were dominated by a moderately intolerant mayfly (*Pseudocloeon*, tolerance value 4.4) and Chironomidae were poorly represented in samples. Wilton (2004) noted that the number of mayfly (Ephemeroptera) taxa in wadeable streams typically exceeds the number of taxa representing the other aquatic insect orders (≤36%), so the dominance of *Pseudocloen* in Dousman Creek should not be considered unsual. Because large numbers of Chironomidae are often associated with disturbance, their low density in Dousman Creek is particularly notable. Taxa richness and EPT richness were lower than expected for this outstanding stream (Iowa Department of Natural Resources, 2010b, Wilton 2004), but the reason for this is unknown. Wilton (2004) reported taxa richness values for reference streams that were generally greater than 20 taxa with about one-third of those being EPT taxa. Several intolerant and uncommon taxa were represented in samples. Water quality and habitat data also suggest this stream is not disturbed. Future invertebrate sampling events at Dousman Creek should include a composited multihabitat sample as described in Wilton (2004). A multihabitat sample in addition to the methodology described in Bowles et al. (2008) will allow investigators to calculate the IOWATER Advanced Benthic Macroinvertebrate Indexing Methods. Using this index to evaluate future benthic collections from Dousman Creek will allow site managers to directly compare its invertebrate community structure relative to Iowa reference streams (Wilton 2004) in addition to assessing long-term trend in this resource.

Literature Cited

- Bowles, D. E., M. H. Williams, H. R. Dodd, L. W. Morrison, J. A. Hinsey, C. E. Ciak, G. A. Rowell, M. D. DeBacker, and J. L. Haack. 2008. Monitoring Protocol for Aquatic Invertebrates of Small Streams in the Heartland Inventory & Monitoring Network. Natural Resource Report NPS/HTLN/NRR—2008/042. National Park Service, Fort Collins, Colorado.
- DeBacker, M. D., C. C. Young (editor), P. Adams., L. Morrison, D. Peitz, G. A. Rowell, M. Williams, and D. Bowles. 2005. Heartland Inventory and Monitoring and Prairie Cluster Prototype Monitoring Program vital signs monitoring plan. U.S. National Park Service, Heartland I&M Network and Prairie Cluster Prototype Monitoring Program, Wilson's Creek National Battlefield, Republic, Missouri.
- Iowa Department of Natural Resources 2010a. Antidegradation, Iowa outstanding waters. http://www.iowadnr.gov/water/standards/antidegradation.html. Accessed: April 2010.
- Iowa Department of Natural Resources. 2010b. Water quality standards. Des Moines, Iowa. http://www.iowadnr.gov/water/standards/criteria.html. Accessed March 8, 2010.
- Wentworth, C. K. 1922.A scale of grade and class terms for clastic sediments. Journal of Geology **30**:377-392.
- Wilton, T. 2004. Biological assessment of Iowa's wadeable streams. Project Report. Iowa Department of Natural Resources, Des Moines, Iowa.

Appendix. Aquatic invertebrate data collected from Dousman Creek, Effigy Mounds National Monument, 2008. FFG= functional feeding group, TV= tolerance value.

Phylum	Class	Order	Family	Genus	Feeding Group	Tolerance Value	1 L	1 M	1 R	2 L	2 M	2 R	3 L	3 M	3 R
Nematomorpha						5					2				1
Annelida	Oligochaeta					8				4	10	5	3		28
	Hirudinea					10								1	
	Hirudinea	Rhynchobdellida	Glossiphoniidae	Glossiphonia		7			1						
Arthropoda	Arachnoida	Hydracarina			Pa, Pr	5.7		2		1	3				1
	Entognatha	Collembola				10			1						3
	Insecta	Ephemeroptera	Baetidae		С	4	13	20			3			30	9
			Baetidae	Pseudocloeon	С	4.4	116	147	138	185	181	114	178	104	85
		Trichoptera	Brachycentridae	Brachycentrus	F	1								1	
			Glossosomatidae	Glossosoma	C, Sc	1.5				1					
			Hydropsychidae		С	4					1				
			Uenoidae	Neophylax	Sc	1.6			1				1		
		Coleoptera	Elmidae	Optioservus	C, Sh	2.7		1		2				1	1
			Dytiscidae	Agabus	Pr	5								1	
	Insecta	Diptera	unknown								1				

Appendix. Continued.

Phylum	Class	Order	Family	Genus	Feeding Group	Tolerance Value	1 L	1 M	1 R	2 L	2 M	2 R	3 L	3 M	3 R
			Chironomidae			6	8	12	14	23	21	19	41	13	39
			Muscidae	Limnophora	Pr	7							1		
			Tipulidae	Dicranota	Pr	0	1	1	1	1	1	1	4	1	5
				Tipula	C, Sh	7.7								1	
			Simuliidae	Simulium	F	4.4	53	101	58	34	81	70	158	74	18
	Crustacea	Amphipoda	Gammaridae	Gammarus	С	6.9	8	3	3	2	2	3	11	2	8
		Isopoda	Asellidae	Lirceus	С	7.7	1	2	3	1		6	10	3	5
Mollusca	Gastropoda	Bassommatophora	Physidae	Physa	Sc	8				1					1
			TOTAL				213	289	220	255	304	220	407	232	204